


CLAIMS

- 
1. In a method which calls for administration of IFN- α , the improvement comprising co-administering an effective amount of an isolated immunostimulatory nucleic acid.
2. The improvement of claim 1, wherein the IFN- α is administered at a dose below the clinically established effective dose for IFN- α alone.
3. The improvement of claim 1, wherein the IFN- α is administered at the maximum tolerated dose for IFN- α in the absence of the nucleic acid.
4. The improvement of claim 1, wherein the IFN- α is administered at least 20 percent below the maximum tolerated dose of IFN- α in the subject.
5. The improvement of claim 1, wherein the IFN- α is administered at least 30 percent below the maximum tolerated dose of IFN- α in the subject.
6. The improvement of claim 1, wherein the IFN- α is administered at least 40 percent below the maximum tolerated dose of IFN- α in the subject.
7. The improvement of claim 1, wherein the IFN- α is administered at least 50 percent below the maximum tolerated dose of IFN- α in the subject.
8. The improvement of claim 1, wherein the immunostimulatory nucleic acid is modified.
9. The improvement of claim 1, wherein the immunostimulatory nucleic acid comprises a backbone with at least one nuclease-resistant internucleotide linkage selected from the group consisting of: phosphorothioate, phosphorodithioate, methylphosphonate, and peptide.

10. The improvement of claim 1, wherein the immunostimulatory nucleic acid comprises at least one nucleotide analog or derivative.

11. The improvement of claim 1, wherein the immunostimulatory nucleic acid is not a
5 palindrome.

12. The improvement of claim 1, wherein the immunostimulatory nucleic acid is a CpG nucleic acid.

10 13. The improvement of claim 1, wherein the immunostimulatory nucleic acid is a non-CpG nucleic acid.

14. The improvement of claim 13, wherein the non-CpG immunostimulatory nucleic acid is a T-rich nucleic acid.

15 15. The improvement of claim 13, wherein the non-CpG immunostimulatory nucleic acid is a poly-G nucleic acid.

16. The improvement of claim 1, wherein the immunostimulatory nucleic acid is any
20 combination of at least two nucleic acids selected from the group consisting of: CpG nucleic acids, T-rich nucleic acids, and poly-G nucleic acids.

17. The improvement of claim 1, wherein the immunostimulatory nucleic acid is between 8 and 100 nucleotides in length.

25 18. The improvement of claim 1, wherein the immunostimulatory nucleic acid is between 12 and 40 nucleotides in length.

30 19. The improvement of claim 1, wherein the immunostimulatory nucleic acid has a sequence selected from the group consisting of

ggGGTCAACGTTGAgggggG
tcgtcgttttgcgttttgcgtt
ggggtcgtcgttttgggggg

ODN 1585 SEQ ID NO:1
ODN 2022 SEQ ID NO:2
ODN 2184 SEQ ID NO:3

002260 " 92727950

Sub
32

tcgtcgttttgtcgttttgggggg
ggggtcgacgtcgagggggg
ggggtcacgatgagggggg
ggGGGACGATCGTCggggggG
5 gggggtcgtacgacgggggg
ggGGGACGATATCGTCggggggG
ggGGGACGACGTCGTCggggggG
ggGGGACGAGCTCGTCggggggG
ggGGGACGTACGTCggggggG
10 ggGGGACGATCGTTGgggggG
ggGGAACGATCGTCggggggG
ggGGGACGATCGTCggggggG
ggGGGACGATCGTCggggggG
ggGGGTCATCGATGAggggggG
15 ggGGTCGTCGACGAggggggG
ggGGTCGTTTGAACGAggggggG
ggGGACGTTTGAACGTggggggG
ggGGAACGACGTCGTTggggggG
ggGGAACGTACGTCggggggG
20 ggGGAACGTACGTACGTTggggggG
ggGGTCACCGGTGAggggggG
ggGGTCGACGTACGTCGAggggggG
ggGGACCGGTACCGGTggggggG
ggGTCGACGTCGAggggggG
25 ggGGTCGACGTCGagggg
ggGGAACGTTAACGTTggggggG
ggGGACGTCGACGTgggggG
ggGGGTCGTTTCGTTggggggG
ggGACGATCGTCGggggggG
30 ggGTCGTCGACGAggggggG
ggTCGTCGACGAGggggggG
ggGGACGATCGTCGggggggG
ggGGTCGACGTCGACGTCGAGggggggG
ggGGACGACGTCGTGggggggG

ODN 2185 SEQ ID NO:4
ODN 2192 SEQ ID NO:5
ODN 2204 SEQ ID NO:6
ODN 2216 SEQ ID NO:7
ODN 2217 SEQ ID NO:8
ODN 2245 SEQ ID NO:9
ODN 2246 SEQ ID NO:10
ODN 2247 SEQ ID NO:11
ODN 2248 SEQ ID NO:12
ODN 2252 SEQ ID NO:13
ODN 2253 SEQ ID NO:14
ODN 2254 SEQ ID NO:15
ODN 2255 SEQ ID NO:16
ODN 2260 SEQ ID NO:17
ODN 2293 SEQ ID NO:18
ODN 2294 SEQ ID NO:19
ODN 2295 SEQ ID NO:20
ODN 2297 SEQ ID NO:21
ODN 2298 SEQ ID NO:22
ODN 2299 SEQ ID NO:23
ODN 2300 SEQ ID NO:24
ODN 2301 SEQ ID NO:25
ODN 2302 SEQ ID NO:26
ODN 2303 SEQ ID NO:27
ODN 2304 SEQ ID NO:28
ODN 2305 SEQ ID NO:29
ODN 2306 SEQ ID NO:30
ODN 2311 SEQ ID NO:31
ODN 2328 SEQ ID NO:32
ODN 2329 SEQ ID NO:33
ODN 2330 SEQ ID NO:34
ODN 2332 SEQ ID NO:35
ODN 2334 SEQ ID NO:36, and
ODN 2336 SEQ ID NO:37,

wherein each lower case letter represents phosphorothioate linkage and each upper case letter indicates phosphodiester linkage.

20. The improvement of claim 1, further comprising co-administering GM-CSF to the subject.

21. The improvement of claim 1, wherein the subject has a condition selected from the group consisting of a proliferative disorder and a viral infection.

22. The improvement of claim 1, wherein the subject has a proliferative disorder selected from the group consisting of: hairy cell leukemia, chronic myelogenous leukemia, cutaneous T-cell leukemia, multiple myeloma, follicular lymphoma, malignant melanoma, squamous cell carcinoma, AIDS-related Kaposi's sarcoma, renal cell carcinoma, prostate carcinoma, bladder cell carcinoma, cervical dysplasia, and colon carcinoma.

23. The improvement of claim 1, wherein the subject has a viral infection selected from the group consisting of: hepatitis B, hepatitis C, condyloma acuminatum, human immunodeficiency virus, herpes, cytomegalovirus, Epstein-Barr virus, and papillomavirus.

24. A method of supplementing IFN- α treatment of a subject comprising administering to a subject in need of IFN- α treatment an effective amount of IFN- α and an isolated immunostimulatory nucleic acid.

25. The method of claim 24, wherein the IFN- α is administered at a dose below the clinically established effective dose for IFN- α alone.

26. The method of claim 24, wherein the IFN- α is administered at the maximum tolerated dose for IFN- α in the absence of the immunostimulatory nucleic acid.

27. The method of claim 24, wherein the IFN- α is administered at least 20 percent below the maximum tolerated dose of IFN- α in the subject.

28. The method of claim 24, wherein the IFN- α is administered at least 30 percent below the maximum tolerated dose of IFN- α in the subject.

29. The method of claim 24, wherein the IFN- α is administered at least 40 percent below the maximum tolerated dose of IFN- α in the subject.

002260" 92700 0964126 092700

133

30. The method of claim 24, wherein the IFN- α is administered at least 50 percent below the maximum tolerated dose of IFN- α in the subject.
31. The method of claim 24, wherein the immunostimulatory nucleic acid is modified.
32. The method of claim 24, wherein the immunostimulatory nucleic acid comprises a backbone with at least one nuclease-resistant internucleotide linkage selected from the group consisting of: phosphorothioate, phosphorodithioate, methylphosphonate, and peptide.
33. The method of claim 24, wherein the immunostimulatory nucleic acid comprises at least one nucleotide analog or derivative.
34. The method of claim 24, wherein the immunostimulatory nucleic acid is not a palindrome.
35. The method of claim 24, wherein the immunostimulatory nucleic acid is a CpG nucleic acid.
36. The method of claim 24, wherein the immunostimulatory nucleic acid is a non-CpG nucleic acid.
37. The method of claim 36, wherein the non-CpG immunostimulatory nucleic acid is a T-rich nucleic acid.
38. The method of claim 36, wherein the non-CpG immunostimulatory nucleic acid is a poly-G nucleic acid.
39. The method of claim 24, wherein the immunostimulatory nucleic acid is any combination of at least two nucleic acids selected from the group consisting of: CpG nucleic acids, T-rich nucleic acids, and poly-G nucleic acids.

40. The method of claim 24, wherein the immunostimulatory nucleic acid is between 8 and 100 nucleotides in length.

41. The method of claim 24, wherein the immunostimulatory nucleic acid is between 12 and 40 nucleotides in length.

42. The method of claim 24, wherein the immunostimulatory nucleic acid has a sequence selected from the group consisting of

ggGGTCAACGTTGAgggggG	ODN 1585	SEQ ID NO:1
tcgtcgttttgcgttttgcgtt	ODN 2022	SEQ ID NO:2
ggggtcgtcgttttggggg	ODN 2184	SEQ ID NO:3
tcgtcgttttgcgttttggggg	ODN 2185	SEQ ID NO:4
ggggtcgacgtcgaggggg	ODN 2192	SEQ ID NO:5
ggggtcacgtcgaggggg	ODN 2204	SEQ ID NO:6
ggGGGACGATCGTCgggggG	ODN 2216	SEQ ID NO:7
gggggtcgtacgacggggg	ODN 2217	SEQ ID NO:8
ggGGGACGATATCGTCgggggG	ODN 2245	SEQ ID NO:9
ggGGGACGACGTCGTCgggggG	ODN 2246	SEQ ID NO:10
ggGGGACGAGCTCGTCgggggG	ODN 2247	SEQ ID NO:11
ggGGGACGTACGTCgggggG	ODN 2248	SEQ ID NO:12
ggGGGACGATCGTTGggggG	ODN 2252	SEQ ID NO:13
ggGGAACGATCGTCgggggG	ODN 2253	SEQ ID NO:14
ggGGGGACGATCGTCgggggG	ODN 2254	SEQ ID NO:15
ggGGGACGATCGTCGgggggG	ODN 2255	SEQ ID NO:16
ggGGGTCATCGATGAgggggG	ODN 2260	SEQ ID NO:17
ggGGTCGTCGACGAgggggG	ODN 2293	SEQ ID NO:18
ggGGTCGTTTCAACGAgggggG	ODN 2294	SEQ ID NO:19
ggGGACGTTTCAACGTgggggG	ODN 2295	SEQ ID NO:20
ggGGAACGACGTCGTTgggggG	ODN 2297	SEQ ID NO:21
ggGGAACGTACGTCgggggG	ODN 2298	SEQ ID NO:22
ggGGAACGTACGTACGTTgggggG	ODN 2299	SEQ ID NO:23
ggGGTCACCGGTGAgggggG	ODN 2300	SEQ ID NO:24
ggGGTCGACGTACGTCGAgggggG	ODN 2301	SEQ ID NO:25
ggGGACCGGTACCGGTgggggG	ODN 2302	SEQ ID NO:26
ggGTCGACGTCGAgggggG	ODN 2303	SEQ ID NO:27
ggGGTCGACGTCGAgggg	ODN 2304	SEQ ID NO:28
ggGGAACGTTAACGTTgggggG	ODN 2305	SEQ ID NO:29
ggGGACGTCGACGTgggggG	ODN 2306	SEQ ID NO:30
ggGGGTCGTTTCGTTgggggG	ODN 2311	SEQ ID NO:31
ggGACGATCGTCGgggggG	ODN 2328	SEQ ID NO:32
ggGTCGTCGACGAggggggG	ODN 2329	SEQ ID NO:33
ggTCGTCGACGAGgggggG	ODN 2330	SEQ ID NO:34
ggGGACGATCGTCGgggggG	ODN 2332	SEQ ID NO:35
ggGGTCGACGTCGACGTCGAGgggggG	ODN 2334	SEQ ID NO:36, and

ggGGACGACGTCGTGgggggG

ODN 2336 SEQ ID NO:37,

wherein each lower case letter represents phosphorothioate linkage and each upper case letter indicates phosphodiester linkage.

43. The method of claim 24, further comprising co-administering GM-CSF to the subject.

44. The method of claim 24, wherein the subject has a condition selected from the group consisting of a proliferative disorder and a viral infection.

45. The method of claim 24, wherein the subject has a proliferative disorder selected from the group consisting of: hairy cell leukemia, chronic myelogenous leukemia, cutaneous T-cell leukemia, multiple myeloma, follicular lymphoma, malignant melanoma, squamous cell carcinoma, AIDS-related Kaposi's sarcoma, renal cell carcinoma, prostate carcinoma, bladder cell carcinoma, cervical dysplasia, and colon carcinoma.

46. The method of claim 24, wherein the subject has a viral infection selected from the group consisting of: hepatitis B, hepatitis C, condyloma acuminatum, human immunodeficiency virus, herpes, cytomegalovirus, Epstein-Barr virus, and papillomavirus.

47. A method of treating a subject to activate interferon-producing cells (IPCs) of the subject comprising
isolating IPCs from a subject in need of such treatment,
culturing the IPCs *in vitro*,
contacting the IPCs *in vitro* with an effective amount of an isolated immunostimulatory nucleic acid, and
returning the contacted IPCs to the subject.

48. The method of claim 47, further comprising contacting the IPCs *in vitro* with a growth factor.

49. The method of claim 47, further comprising contacting the IPCs *in vitro* with IL-3.

50. The method of claim 47, further comprising contacting the IPCs *in vitro* with GM-CSF.
51. The method of claim 47, wherein the IPCs are cultured *in vitro* in the absence of IL-3.
52. The method of claim 47, wherein the IPCs are cultured *in vitro* in the absence of GM-CSF.
53. The method of claim 47, wherein the immunostimulatory nucleic acid is modified.
54. The method of claim 47, wherein the immunostimulatory nucleic acid comprises a backbone with at least one nuclease-resistant internucleotide linkage selected from the group consisting of: phosphorothioate, phosphorodithioate, methylphosphonate, and peptide.
55. The method of claim 47, wherein the immunostimulatory nucleic acid comprises at least one nucleotide analog or derivative.
56. The method of claim 47, wherein the immunostimulatory nucleic acid is not a palindrome.
57. The method of claim 47, wherein the immunostimulatory nucleic acid is a CpG nucleic acid.
58. The method of claim 47, wherein the immunostimulatory nucleic acid is a non-CpG nucleic acid.
59. The method of claim 58, wherein the non-CpG immunostimulatory nucleic acid is a T-rich nucleic acid.
60. The method of claim 58, wherein the non-CpG immunostimulatory nucleic acid is a poly-G nucleic acid.

61. The method of claim 47, wherein the immunostimulatory nucleic acid is any combination of at least two nucleic acids selected from the group consisting of: CpG nucleic acids, T-rich nucleic acids, and poly-G nucleic acids.

62. The method of claim 47, wherein the immunostimulatory nucleic acid is between 8 and 100 nucleotides in length.

63. The method of claim 47, wherein the immunostimulatory nucleic acid is between 12 and 40 nucleotides in length.

64. The method of claim 47, wherein the immunostimulatory nucleic acid has a sequence selected from the group consisting of

ggGGTCAACGTTGAgggggG	ODN 1585	SEQ ID NO:1
tcgtcgtttgcgttttgcgtt	ODN 2022	SEQ ID NO:2
ggggtcgctgttttgggggg	ODN 2184	SEQ ID NO:3
tcgtcgtttgcgttttgggggg	ODN 2185	SEQ ID NO:4
ggggtcgacgtcgagggggg	ODN 2192	SEQ ID NO:5
ggggtcatcgatgagggggg	ODN 2204	SEQ ID NO:6
ggGGGACGATCGTCgggggG	ODN 2216	SEQ ID NO:7
gggggtcgtagcagggggg	ODN 2217	SEQ ID NO:8
ggGGGACGATATCGTCgggggG	ODN 2245	SEQ ID NO:9
ggGGGACGACGTCGTCgggggG	ODN 2246	SEQ ID NO:10
ggGGGACGAGCTCGTCgggggG	ODN 2247	SEQ ID NO:11
ggGGGACGTACGTCgggggG	ODN 2248	SEQ ID NO:12
ggGGGACGATCGTTGgggggG	ODN 2252	SEQ ID NO:13
ggGGAACGATCGTCgggggG	ODN 2253	SEQ ID NO:14
ggGGGACGATCGTCgggggG	ODN 2254	SEQ ID NO:15
ggGGGACGATCGTCgggggG	ODN 2255	SEQ ID NO:16
ggGGGTCATCGATGAgggggG	ODN 2260	SEQ ID NO:17
ggGGTCGTCGACGAgggggG	ODN 2293	SEQ ID NO:18
ggGGTCGTTTGAACGAgggggG	ODN 2294	SEQ ID NO:19
ggGGACGTTTGAACGTgggggG	ODN 2295	SEQ ID NO:20
ggGGAACGACGTCGTTgggggG	ODN 2297	SEQ ID NO:21
ggGGAACGTACGTCgggggG	ODN 2298	SEQ ID NO:22
ggGGAACGTACGTACGTTgggggG	ODN 2299	SEQ ID NO:23
ggGGTCACCGGTGAgggggG	ODN 2300	SEQ ID NO:24
ggGGTCGACGTACGTCGAgggggG	ODN 2301	SEQ ID NO:25
ggGGACCGGTACCGGTgggggG	ODN 2302	SEQ ID NO:26
ggGTCGACGTCGAgggggG	ODN 2303	SEQ ID NO:27
ggGGTCGACGTCGAgggg	ODN 2304	SEQ ID NO:28
ggGGAACGTTAACGTTgggggG	ODN 2305	SEQ ID NO:29

002260 92126 092700

0000079628

10

- 15

20

25

30

ggGGGACGATATCGTCggggggG	ODN 2245	SEQ ID NO:9
ggGGGACGACGTCGTCggggggG	ODN 2246	SEQ ID NO:10
ggGGGACGAGCTCGTCggggggG	ODN 2247	SEQ ID NO:11
ggGGGACGTACGTCggggggG	ODN 2248	SEQ ID NO:12
5 ggGGGACGATCGTTGgggggG	ODN 2252	SEQ ID NO:13
ggGGAACGATCGTCggggggG	ODN 2253	SEQ ID NO:14
ggGGGGACGATCGTCggggggG	ODN 2254	SEQ ID NO:15
ggGGGACGATCGTCGggggggG	ODN 2255	SEQ ID NO:16
ggGGGTCATCGATGAggggggG	ODN 2260	SEQ ID NO:17
10 ggGGTCGTCGACGAggggggG	ODN 2293	SEQ ID NO:18
ggGGTCGTTCTGAACGAggggggG	ODN 2294	SEQ ID NO:19
ggGGACGTTCTGAACGTggggggG	ODN 2295	SEQ ID NO:20
ggGGAACGACGTCGTTggggggG	ODN 2297	SEQ ID NO:21
ggGGAACGTACGTCggggggG	ODN 2298	SEQ ID NO:22
15 ggGGAACGTACGTACGTTggggggG	ODN 2299	SEQ ID NO:23
ggGGTCACCGGTGAggggggG	ODN 2300	SEQ ID NO:24
ggGGTCGACGTACGTCGAggggggG	ODN 2301	SEQ ID NO:25
ggGGACCGGTACCGGTggggggG	ODN 2302	SEQ ID NO:26
ggGTCGACGTCGAggggggG	ODN 2303	SEQ ID NO:27
20 ggGGTCGACGTCGAggggg	ODN 2304	SEQ ID NO:28
ggGGAACGTTAACGTTggggggG	ODN 2305	SEQ ID NO:29
ggGGACGTCGACGTggggggG	ODN 2306	SEQ ID NO:30
ggGGGTCGTTCTGTTggggggG	ODN 2311	SEQ ID NO:31
ggGACGATCGTCGggggggG	ODN 2328	SEQ ID NO:32
25 ggGTCGTCGACGAgggggggG	ODN 2329	SEQ ID NO:33
ggTCGTCGACGAGggggggG	ODN 2330	SEQ ID NO:34
ggGGACGATCGTCGggggggG	ODN 2332	SEQ ID NO:35
ggGGTCGACGTCGACGTCGAGggggggG	ODN 2334	SEQ ID NO:36, and
ggGGACGACGTCGTGggggggG	ODN 2336	SEQ ID NO:37,

30 wherein each lower case letter represents phosphorothioate linkage and each upper case letter indicates phosphodiester linkage.

79. The method of claim 65, wherein the subject has a condition selected from the group consisting of a proliferative disorder and a viral infection.

80. The method of claim 65, wherein the subject has a proliferative disorder selected from the group consisting of: hairy cell leukemia, chronic myelogenous leukemia, cutaneous T-cell leukemia, multiple myeloma, follicular lymphoma, malignant melanoma, squamous cell carcinoma, AIDS-related Kaposi's sarcoma, renal cell carcinoma, prostate carcinoma, bladder cell carcinoma, cervical dysplasia, and colon carcinoma.

81. The method of claim 65, wherein the subject has a viral infection selected from the group consisting of: hepatitis B, hepatitis C, condyloma acuminatum, human immunodeficiency virus, herpes, cytomegalovirus, Epstein-Barr virus, and papillomavirus.

5
Sub 805
82. A method of decreasing a dose of IFN- α effective for treating a subject, comprising: administering to a subject in need of treatment with IFN- α a pharmaceutical composition comprising IFN- α , and coadministering to the subject in need of such treatment a pharmaceutical composition comprising an immunostimulatory nucleic acid in an amount which, together with the administered IFN- α , is an effective IFN- α treatment, and, wherein the amount of administered IFN- α is less than an amount of IFN- α required in the absence of coadministering the immunostimulatory nucleic acid.

10
15 83. The method of claim 82, wherein the amount of administered IFN- α is at least 20 percent below the amount of IFN- α required in the absence of coadministering the immunostimulatory nucleic acid.

20 84. The method of claim 82, wherein the amount of administered IFN- α is at least 30 percent below the amount of IFN- α required in the absence of coadministering the immunostimulatory nucleic acid.

25 85. The method of claim 82, wherein the amount of administered IFN- α is at least 40 percent below the amount of IFN- α required in the absence of coadministering the immunostimulatory nucleic acid.

30 86. The method of claim 82, wherein the amount of administered IFN- α is at least 50 percent below the amount of IFN- α required in the absence of coadministering the immunostimulatory nucleic acid.

00672126.092700

87. The method of claim 82, wherein the pharmaceutical composition comprising an immunostimulatory nucleic acid is administered locally.
88. The method of claim 82, wherein the immunostimulatory nucleic acid is modified.
89. The method of claim 82, wherein the immunostimulatory nucleic acid comprises a backbone with at least one nuclease-resistant internucleotide linkage selected from the group consisting of: phosphorothioate, phosphorodithioate, methylphosphonate, and peptide.
90. The method of claim 82, wherein the immunostimulatory nucleic acid comprises at least one nucleotide analog or derivative.
91. The method of claim 82, wherein the immunostimulatory nucleic acid is not a palindrome.
92. The method of claim 82, wherein the immunostimulatory nucleic acid is a CpG nucleic acid.
93. The method of claim 82, wherein the immunostimulatory nucleic acid is a non-CpG nucleic acid.
94. The method of claim 93, wherein the non-CpG immunostimulatory nucleic acid is a T-rich nucleic acid.
95. The method of claim 93, wherein the non-CpG immunostimulatory nucleic acid is a poly-G nucleic acid.
96. The method of claim 82, wherein the immunostimulatory nucleic acid is any combination of at least two nucleic acids selected from the group consisting of: CpG nucleic acids, T-rich nucleic acids, and poly-G nucleic acids.

97. The method of claim 82, wherein the immunostimulatory nucleic acid is between 8 and 100 nucleotides in length.

98. The method of claim 82, wherein the immunostimulatory nucleic acid is between 12 and 40 nucleotides in length.

99. The method of claim 82, wherein the immunostimulatory nucleic acid has a sequence selected from the group consisting of

ggGGTCAACGTTGAggggggG	ODN 1585	SEQ ID NO:1
tcgtcgttttgcgttttgcgtt	ODN 2022	SEQ ID NO:2
ggggtcgtcgttttgggggg	ODN 2184	SEQ ID NO:3
tcgtcgttttgcgttttgggggg	ODN 2185	SEQ ID NO:4
ggggtcgacgtcgagggggg	ODN 2192	SEQ ID NO:5
ggggtcacgtcgagggggg	ODN 2204	SEQ ID NO:6
ggGGGACGATCGTCggggggG	ODN 2216	SEQ ID NO:7
gggggtcgtagcagggggg	ODN 2217	SEQ ID NO:8
ggGGGACGATATCGTCggggggG	ODN 2245	SEQ ID NO:9
ggGGGACGACGTCGTCggggggG	ODN 2246	SEQ ID NO:10
ggGGGACGAGCTCGTCggggggG	ODN 2247	SEQ ID NO:11
ggGGGACGTACGTCggggggG	ODN 2248	SEQ ID NO:12
ggGGGACGATCGTTGgggggG	ODN 2252	SEQ ID NO:13
ggGGAACGATCGTCggggggG	ODN 2253	SEQ ID NO:14
ggGGGACGATCGTCggggggG	ODN 2254	SEQ ID NO:15
ggGGGACGATCGTCGggggggG	ODN 2255	SEQ ID NO:16
ggGGGTCATCGATGAggggggG	ODN 2260	SEQ ID NO:17
ggGGTCGTCGACGAggggggG	ODN 2293	SEQ ID NO:18
ggGGTCGTTTCAACGAggggggG	ODN 2294	SEQ ID NO:19
ggGGACGTTTCAACGTggggggG	ODN 2295	SEQ ID NO:20
ggGGAACGACGTCGTTggggggG	ODN 2297	SEQ ID NO:21
ggGGAACGTACGTCggggggG	ODN 2298	SEQ ID NO:22
ggGGAACGTACGTACGTTggggggG	ODN 2299	SEQ ID NO:23
ggGGTCACCGGTGAggggggG	ODN 2300	SEQ ID NO:24
ggGGTCGACGTACGTCGAggggggG	ODN 2301	SEQ ID NO:25
ggGGACCGGTACCGGTggggggG	ODN 2302	SEQ ID NO:26
ggGGTCGACGTGAggggggG	ODN 2303	SEQ ID NO:27
ggGGTCGACGTGAggggg	ODN 2304	SEQ ID NO:28
ggGGAACGTTAACGTTggggggG	ODN 2305	SEQ ID NO:29
ggGGACGTGACGTggggggG	ODN 2306	SEQ ID NO:30
ggGGGTCGTTTCGTTggggggG	ODN 2311	SEQ ID NO:31
ggGACGATCGTCGggggggG	ODN 2328	SEQ ID NO:32
ggGTCGTCGACGAgggggggG	ODN 2329	SEQ ID NO:33
ggTCGTCGACGAggggggG	ODN 2330	SEQ ID NO:34
ggGGACGATCGTCGggggggG	ODN 2332	SEQ ID NO:35
ggGGTCGACGTCGACGTCGAGggggggG	ODN 2334	SEQ ID NO:36, and

ggGACGACGTCGTGgggggG

ODN 2336 SEQ ID NO:37,

wherein each lower case letter represents phosphorothioate linkage and each upper case letter indicates phosphodiester linkage.

100. The method of claim 82, wherein the subject has a condition selected from the group consisting of a proliferative disorder and a viral infection.

101. The method of claim 82, wherein the subject has a proliferative disorder selected from the group consisting of: hairy cell leukemia, chronic myelogenous leukemia, cutaneous T-cell leukemia, multiple myeloma, follicular lymphoma, malignant melanoma, squamous cell carcinoma, AIDS-related Kaposi's sarcoma, renal cell carcinoma, prostate carcinoma, bladder cell carcinoma, cervical dysplasia, and colon carcinoma.

102. The method of claim 82, wherein the subject has a viral infection selected from the group consisting of: hepatitis B, hepatitis C, condyloma acuminatum, human immunodeficiency virus, herpes, cytomegalovirus, Epstein-Barr virus, and papillomavirus.

103. A method of preventing an IFN- α treatment-related side effect in a subject receiving or in need of treatment with IFN- α , comprising
administering to a subject in need of treatment with IFN- α a pharmaceutical composition comprising IFN- α , and
coadministering to the subject in need of such treatment a pharmaceutical composition comprising an immunostimulatory nucleic acid in an amount which, together with the administered IFN- α , is an effective IFN- α treatment, and, wherein an IFN- α treatment-related side effect is reduced in comparison to the side effect when IFN- α is administered in the absence of coadministering the immunostimulatory nucleic acid.

104. The method of claim 103, wherein the pharmaceutical composition comprising an immunostimulatory nucleic acid is administered locally.

105. The method of claim 103, wherein the IFN- α treatment-related side effect is systemic.

002260.92T27960

20
Sub B6

106. The method of claim 103, wherein the IFN- α treatment-related side effect is selected from the group consisting of flu-like syndrome, fever, headache, chills, myalgia, fatigue, anorexia, nausea, vomiting, diarrhea, and depression.

5

107. The method of claim 103, wherein the immunostimulatory nucleic acid is modified.

108. The method of claim 103, wherein the immunostimulatory nucleic acid comprises a backbone with at least one nuclease-resistant internucleotide linkage selected from the group consisting of: phosphorothioate, phosphorodithioate, methylphosphonate, and peptide.

10

109. The method of claim 103, wherein the immunostimulatory nucleic acid comprises at least one nucleotide analog or derivative.

15

110. The method of claim 103, wherein the immunostimulatory nucleic acid is not a palindrome.

111. The method of claim 103, wherein the immunostimulatory nucleic acid is a CpG nucleic acid.

20

112. The method of claim 103, wherein the immunostimulatory nucleic acid is a non-CpG nucleic acid.

25

113. The method of claim 112, wherein the non-CpG immunostimulatory nucleic acid is a T-rich nucleic acid.

114. The method of claim 112, wherein the non-CpG immunostimulatory nucleic acid is a poly-G nucleic acid.

30

007260 9272760

115. The method of claim 103, wherein the immunostimulatory nucleic acid is any combination of at least two nucleic acids selected from the group consisting of: CpG nucleic acids, T-rich nucleic acids, and poly-G nucleic acids.

116. The method of claim 103, wherein the immunostimulatory nucleic acid is between 8 and 100 nucleotides in length.

117. The method of claim 103, wherein the immunostimulatory nucleic acid is between 12 and 40 nucleotides in length.

118. The method of claim 103, wherein the immunostimulatory nucleic acid has a sequence selected from the group consisting of

ggGGTCAACGTTGAgggggG	ODN 1585	SEQ ID NO:1
tcgtcgttttgcgttttgcgtt	ODN 2022	SEQ ID NO:2
ggggtcgtcgttttggggggg	ODN 2184	SEQ ID NO:3
tcgtcgttttgcgttttggggggg	ODN 2185	SEQ ID NO:4
ggggtcgacgtcgagggggg	ODN 2192	SEQ ID NO:5
ggggtcacgtcgagggggg	ODN 2204	SEQ ID NO:6
ggGGGACGATCGTCggggggG	ODN 2216	SEQ ID NO:7
gggggtcgtacgacgggggg	ODN 2217	SEQ ID NO:8
ggGGGACGATATCGTCggggggG	ODN 2245	SEQ ID NO:9
ggGGGACGACGTCGTCggggggG	ODN 2246	SEQ ID NO:10
ggGGGACGAGCTCGTCggggggG	ODN 2247	SEQ ID NO:11
ggGGGACGTACGTCggggggG	ODN 2248	SEQ ID NO:12
ggGGGACGATCGTTGggggggG	ODN 2252	SEQ ID NO:13
ggGGAACGATCGTCggggggG	ODN 2253	SEQ ID NO:14
ggGGGGACGATCGTCggggggG	ODN 2254	SEQ ID NO:15
ggGGGACGATCGTCGggggggG	ODN 2255	SEQ ID NO:16
ggGGGTCATCGATGAgggggG	ODN 2260	SEQ ID NO:17
ggGGTCGTCGACGAgggggG	ODN 2293	SEQ ID NO:18
ggGGTCGTTTCAACGAgggggG	ODN 2294	SEQ ID NO:19
ggGGACGTTTCAACGTggggggG	ODN 2295	SEQ ID NO:20
ggGGAACGACGTCGTTggggggG	ODN 2297	SEQ ID NO:21
ggGGAACGTACGTCggggggG	ODN 2298	SEQ ID NO:22
ggGGAACGTACGTACGTTggggggG	ODN 2299	SEQ ID NO:23
ggGGTCACCGGTGAgggggG	ODN 2300	SEQ ID NO:24
ggGGTCGACGTACGTCGAgggggG	ODN 2301	SEQ ID NO:25
ggGGACCGGTACCGGTggggggG	ODN 2302	SEQ ID NO:26
ggGTCGACGTCGAgggggG	ODN 2303	SEQ ID NO:27
ggGGTCGACGTCGAggggg	ODN 2304	SEQ ID NO:28
ggGGAACGTTAACGTTggggggG	ODN 2305	SEQ ID NO:29
ggGGACGTCGACGTgggggG	ODN 2306	SEQ ID NO:30

004260" 9272960

ggGGGTCGTTCGTTgggggG	ODN 2311	SEQ ID NO:31
ggGACGATCGTCGgggggG	ODN 2328	SEQ ID NO:32
ggGTCGTCGACGAggggggG	ODN 2329	SEQ ID NO:33
ggTCGTCGACGAGgggggG	ODN 2330	SEQ ID NO:34
5 ggGGACGATCGTCGgggggG	ODN 2332	SEQ ID NO:35
ggGGTTCGACGTCGACGTCGAGgggggG	ODN 2334	SEQ ID NO:36, and
ggGGACGACGTCGTGgggggG	ODN 2336	SEQ ID NO:37,

wherein each lower case letter represents phosphorothioate linkage and each upper case letter indicates phosphodiester linkage.

119. The method of claim 103, wherein the subject has a condition selected from the group consisting of a proliferative disorder and a viral infection.

120. The method of claim 103, wherein the subject has a proliferative disorder selected from the group consisting of: hairy cell leukemia, chronic myelogenous leukemia, cutaneous T-cell leukemia, multiple myeloma, follicular lymphoma, malignant melanoma, squamous cell carcinoma, AIDS-related Kaposi's sarcoma, renal cell carcinoma, prostate carcinoma, bladder cell carcinoma, cervical dysplasia, and colon carcinoma.

121. The method of claim 103, wherein the subject has a viral infection selected from the group consisting of: hepatitis B, hepatitis C, condyloma acuminatum, human immunodeficiency virus, herpes, cytomegalovirus, Epstein-Barr virus, and papillomavirus.

122. A method of enhancing efficacy of IFN- α treatment in a subject in need of such treatment, comprising

administering to a subject in need of such treatment an amount of a pharmaceutical composition comprising IFN- α effective for treating a condition of the subject;

isolating natural interferon-producing cells (IPCs) from a donor;

contacting the isolated IPCs *ex vivo* with an amount of a pharmaceutical composition comprising an immunostimulatory nucleic acid effective for inducing the IPCs to release IFN- α ; and

administering the contacted cells to the subject.

123. The method of claim 122, wherein the donor is the subject.

124. The method of claim 122 further comprising contacting the isolated IPCs with an
5 antigen.

125. The method of claim 122, wherein the administering the contacted cells comprises
local injection.

10 126. The method of claim 125, wherein the local injection is via a blood vessel supplying a
target tissue.

127. The method of claim 126, wherein the blood vessel is selected from the group
consisting of a hepatic artery, a portal vein, a celiac artery, and a splenic artery.

15 128. The method of claim 122, wherein the immunostimulatory nucleic acid is modified.

129. The method of claim 122, wherein the immunostimulatory nucleic acid comprises a
backbone with at least one nuclease-resistant internucleotide linkage selected from the
20 group consisting of: phosphorothioate, phosphorodithioate, methylphosphonate, and
peptide.

130. The method of claim 122, wherein the immunostimulatory nucleic acid comprises at
least one nucleotide analog or derivative.

25 131. The method of claim 122, wherein the immunostimulatory nucleic acid is not a
palindrome.

132. The method of claim 122, wherein the immunostimulatory nucleic acid is a CpG
30 nucleic acid.

007260-092700

133. The method of claim 122, wherein the immunostimulatory nucleic acid is a non-CpG nucleic acid.

134. The method of claim 133, wherein the non-CpG immunostimulatory nucleic acid is a T-rich nucleic acid.

135. The method of claim 133, wherein the non-CpG immunostimulatory nucleic acid is a poly-G nucleic acid.

136. The method of claim 122, wherein the immunostimulatory nucleic acid is any combination of at least two nucleic acids selected from the group consisting of: CpG nucleic acids, T-rich nucleic acids, and poly-G nucleic acids.

137. The method of claim 122, wherein the immunostimulatory nucleic acid is between 8 and 100 nucleotides in length.

138. The method of claim 122, wherein the immunostimulatory nucleic acid is between 12 and 40 nucleotides in length.

139. The method of claim 122, wherein the immunostimulatory nucleic acid has a sequence selected from the group consisting of

ggGGTCAACGTTGAgggggG

ODN 1585 SEQ ID NO:1

tcgtcgttttgcgttttgcgtt

ODN 2022 SEQ ID NO:2

ggggtcgtcgttttgggggg

ODN 2184 SEQ ID NO:3

tcgtcgttttgcgttttgggggg

ODN 2185 SEQ ID NO:4

ggggtcgacgtcgagggggg

ODN 2192 SEQ ID NO:5

ggggtcacgtcgagggggg

ODN 2204 SEQ ID NO:6

ggGGGACGATCGTCgggggG

ODN 2216 SEQ ID NO:7

gggggtcgtacgacggggg

ODN 2217 SEQ ID NO:8

ggGGGACGATATCGTCgggggG

ODN 2245 SEQ ID NO:9

ggGGGACGACGTCGTCgggggG

ODN 2246 SEQ ID NO:10

ggGGGACGAGCTCGTCgggggG

ODN 2247 SEQ ID NO:11

ggGGGACGTACGTCgggggG

ODN 2248 SEQ ID NO:12

ggGGGACGATCGTTGgggggG

ODN 2252 SEQ ID NO:13

ggGGAACGATCGTCgggggG

ODN 2253 SEQ ID NO:14

ggGGGACGATCGTCgggggG

ODN 2254 SEQ ID NO:15

ggGGGACGATCGTCGgggggG

ODN 2255 SEQ ID NO:16

ggGGGTCATCGATGAgggggG

ODN 2260 SEQ ID NO:17

ggGGTCGTCGACGAggggggG	ODN 2293	SEQ ID NO:18
ggGGTCGTTCTGAACGAggggggG	ODN 2294	SEQ ID NO:19
ggGGACGTTCTGAACGTggggggG	ODN 2295	SEQ ID NO:20
ggGGAACGACGTCGTTggggggG	ODN 2297	SEQ ID NO:21
5 ggGGAACGTACGTCggggggG	ODN 2298	SEQ ID NO:22
ggGGAACGTACGTACGTTggggggG	ODN 2299	SEQ ID NO:23
ggGGTCACCGGTGAggggggG	ODN 2300	SEQ ID NO:24
ggGGTCGACGTACGTCGAggggggG	ODN 2301	SEQ ID NO:25
ggGGACCGGTACCGGTggggggG	ODN 2302	SEQ ID NO:26
10 ggGTCGACGTCGAggggggG	ODN 2303	SEQ ID NO:27
ggGGTCGACGTCGAgggggG	ODN 2304	SEQ ID NO:28
ggGGAACGTTAACGTTggggggG	ODN 2305	SEQ ID NO:29
ggGGACGTCGACGTggggggG	ODN 2306	SEQ ID NO:30
ggGGGTCGTTCTGTTggggggG	ODN 2311	SEQ ID NO:31
15 ggGACGATCGTCGggggggG	ODN 2328	SEQ ID NO:32
ggGTCGTCGACGAgggggggG	ODN 2329	SEQ ID NO:33
ggTCGTCGACGAGggggggG	ODN 2330	SEQ ID NO:34
ggGGACGATCGTCGggggggG	ODN 2332	SEQ ID NO:35
ggGGTCGACGTCGACGTCGAGggggggG	ODN 2334	SEQ ID NO:36, and
20 ggGGACGACGTCGTGggggggG	ODN 2336	SEQ ID NO:37,

wherein each lower case letter represents phosphorothioate linkage and each upper case letter indicates phosphodiester linkage.

140. The method of claim 122, wherein the subject has a condition selected from the group consisting of a proliferative disorder and a viral infection.

141. The method of claim 122, wherein the subject has a proliferative disorder selected from the group consisting of: hairy cell leukemia, chronic myelogenous leukemia, cutaneous T-cell leukemia, multiple myeloma, follicular lymphoma, malignant melanoma, squamous cell carcinoma, AIDS-related Kaposi's sarcoma, renal cell carcinoma, prostate carcinoma, bladder cell carcinoma, cervical dysplasia, and colon carcinoma.

142. The method of claim 122, wherein the subject has a viral infection selected from the group consisting of: hepatitis B, hepatitis C, condyloma acuminatum, human immunodeficiency virus, herpes, cytomegalovirus, Epstein-Barr virus, and papillomavirus.

143. A method of supporting survival of natural interferon-producing cells (IPCs) *in vitro*, comprising
- isolating IPCs from a subject;
 - culturing the IPCs in a sterile medium suitable for tissue culture; and
 - contacting the IPCs *in vitro* with an amount of immunostimulatory nucleic acid effective to support the growth of the IPCs in the absence of interleukin 3 (IL-3).
144. The method of claim 143, wherein the IPCs are precursor type 2 dendritic cells (pDC2s).
145. The method of claim 143, wherein the IPCs are cultured in the absence of IL-3.
146. The method of claim 143, wherein the IPCs are cultured in the absence of GM-CSF.
147. The method of claim 143, wherein the immunostimulatory nucleic acid is modified.
148. The method of claim 143, wherein the immunostimulatory nucleic acid comprises a backbone with at least one nuclease-resistant internucleotide linkage selected from the group consisting of: phosphorothioate, phosphorodithioate, methylphosphonate, and peptide.
149. The method of claim 143, wherein the immunostimulatory nucleic acid comprises at least one nucleotide analog or derivative.
150. The method of claim 143, wherein the immunostimulatory nucleic acid is not a palindrome.
151. The method of claim 143, wherein the immunostimulatory nucleic acid is a CpG nucleic acid.
152. The method of claim 143, wherein the immunostimulatory nucleic acid is a non-CpG nucleic acid.

153. The method of claim 152, wherein the non-CpG immunostimulatory nucleic acid is a T-rich nucleic acid.

5 154. The method of claim 152, wherein the non-CpG immunostimulatory nucleic acid is a poly-G nucleic acid.

10 155. The method of claim 143, wherein the immunostimulatory nucleic acid is any combination of at least two nucleic acids selected from the group consisting of: CpG nucleic acids, T-rich nucleic acids, and poly-G nucleic acids.

156. The method of claim 143, wherein the immunostimulatory nucleic acid is between 8 and 100 nucleotides in length.

15 157. The method of claim 143, wherein the immunostimulatory nucleic acid is between 12 and 40 nucleotides in length.

158. The method of claim 143, wherein the immunostimulatory nucleic acid has a sequence selected from the group consisting of

20 ggGGTCAACGTTGAgggggG
tcgtcgtttgtcgtttgtcgtt
ggggtcgtcgttttgggggg
tcgtcgtttgtcgttttgggggg
ggggtcgacgtcgagggggg
25 ggggtcatcgatgagggggg
ggGGGACGATCGTCggggggG
gggggtcgtacgacgggggg
ggGGGACGATATCGTCggggggG
ggGGGACGACGTCGTCggggggG
30 ggGGGACGAGCTCGTCggggggG
ggGGGACGTACGTCggggggG
ggGGGACGATCGTTGgggggG
ggGGAACGATCGTCggggggG
ggGGGACGATCGTCggggggG
35 ggGGGACGATCGTCGggggggG
ggGGGTCATCGATGAgggggG
ggGGGTCGTCGACGAgggggG
ggGGGTCGTTTGAACGAgggggG
ggGGGACGTTTGAACGTggggggG

ODN 1585	SEQ ID NO:1
ODN 2022	SEQ ID NO:2
ODN 2184	SEQ ID NO:3
ODN 2185	SEQ ID NO:4
ODN 2192	SEQ ID NO:5
ODN 2204	SEQ ID NO:6
ODN 2216	SEQ ID NO:7
ODN 2217	SEQ ID NO:8
ODN 2245	SEQ ID NO:9
ODN 2246	SEQ ID NO:10
ODN 2247	SEQ ID NO:11
ODN 2248	SEQ ID NO:12
ODN 2252	SEQ ID NO:13
ODN 2253	SEQ ID NO:14
ODN 2254	SEQ ID NO:15
ODN 2255	SEQ ID NO:16
ODN 2260	SEQ ID NO:17
ODN 2293	SEQ ID NO:18
ODN 2294	SEQ ID NO:19
ODN 2295	SEQ ID NO:20

ggGGAACGACGTCGTTggggggG	ODN 2297	SEQ ID NO:21
ggGGAACGTACGTCggggggG	ODN 2298	SEQ ID NO:22
ggGGAACGTACGTACGTTggggggG	ODN 2299	SEQ ID NO:23
ggGGTCACCGGTGAggggggG	ODN 2300	SEQ ID NO:24
5 ggGGTCGACGTACGTCGAggggggG	ODN 2301	SEQ ID NO:25
ggGGACCGGTACCGGTggggggG	ODN 2302	SEQ ID NO:26
ggGTCGACGTCGAggggggG	ODN 2303	SEQ ID NO:27
ggGGTCGACGTCGagggg	ODN 2304	SEQ ID NO:28
ggGGAACGTTAACGTTggggggG	ODN 2305	SEQ ID NO:29
10 ggGGACGTCGACGTgggggG	ODN 2306	SEQ ID NO:30
ggGGGTCGTTTCGTTggggggG	ODN 2311	SEQ ID NO:31
ggGACGATCGTCGggggggG	ODN 2328	SEQ ID NO:32
ggGTCGTCGACGAgggggggG	ODN 2329	SEQ ID NO:33
ggTCGTCGACGAGggggggG	ODN 2330	SEQ ID NO:34
15 ggGGACGATCGTCGggggggG	ODN 2332	SEQ ID NO:35
ggGGTCGACGTCGACGTCGAGggggggG	ODN 2334	SEQ ID NO:36, and
ggGGACGACGTCGTGggggggG	ODN 2336	SEQ ID NO:37,

wherein each lower case letter represents phosphorothioate linkage and each upper case letter indicates phosphodiester linkage.

159. A method of stimulating isolated interferon-producing cells (IPCs) *in vitro*, comprising

- isolating IPCs from a subject;
- culturing the IPCs in a sterile medium suitable for tissue culture; and
- 25 contacting the IPCs *in vitro* with an amount of immunostimulatory nucleic acid effective to induce secretion of at least one type I interferon.

160. The method of claim 159, wherein the IPCs are precursor type 2 dendritic cells (pDC2s).

161. The method of claim 159, wherein the type I interferon is an IFN- α .

162. The method of claim 159, wherein the IPCs are cultured in the absence of IL-3.

35 163. The method of claim 159, wherein the IPCs are cultured in the absence of GM-CSF.

164. The method of claim 159, wherein the immunostimulatory nucleic acid is modified.

165. The method of claim 159, wherein the immunostimulatory nucleic acid comprises a backbone with at least one nuclease-resistant internucleotide linkage selected from the group consisting of: phosphorothioate, phosphorodithioate, methylphosphonate, and peptide.

166. The method of claim 159, wherein the immunostimulatory nucleic acid comprises at least one nucleotide analog or derivative.

167. The method of claim 159, wherein the immunostimulatory nucleic acid is not a palindrome.

168. The method of claim 159, wherein the immunostimulatory nucleic acid is a CpG nucleic acid.

169. The method of claim 159, wherein the immunostimulatory nucleic acid is a non-CpG nucleic acid.

170. The method of claim 169, wherein the non-CpG immunostimulatory nucleic acid is a T-rich nucleic acid.

171. The method of claim 169, wherein the non-CpG immunostimulatory nucleic acid is a poly-G nucleic acid.

172. The method of claim 159, wherein the immunostimulatory nucleic acid is any combination of at least two nucleic acids selected from the group consisting of: CpG nucleic acids, T-rich nucleic acids, and poly-G nucleic acids.

173. The method of claim 159, wherein the immunostimulatory nucleic acid is between 8 and 100 nucleotides in length.

174. The method of claim 159, wherein the immunostimulatory nucleic acid is between 12 and 40 nucleotides in length.

002260" 92127960

ggGGTCAACGTTGAggggggG

ODN 2022 SEQ ID NO:2

ODN 2184 SEQ ID NO:3

ODN 2185 SEQ ID NO:4

ODN 2192 SEO ID NO:5

ODN 2204 / SEO ID NO:6

ODN 2216 ✓ SEQ ID NO:7

ODN 2217 SEQ ID NO:8

ODN 2245 SEQ ID NO:9

ODN 2246 SEQ ID NO:1

QDN 2247 SEQ ID NO:1

ODN 2248 SEQ ID NO:1

ODN 2248 SEQ ID NO:1
ODN 2252 SEQ ID NO:1

ODN 2253 SEQ ID NO:1

ODN 2254 SEQ ID NO:1

ODN 2254 SEQ ID NO:1
ODN 2255 SEQ ID NO:1

ODN 2255 SEQ ID NO:1
ODN 2260 SEQ ID NO:1

ODN 2260 SEQ ID NO:1
ODN 2293 SEQ ID NO:1

ODN 2293 SEQ ID NO:1
ODN 2294 SEQ ID NO:1

ODN 2294 SEQ ID NO:1
ODN 2295 SEQ ID NO:2

ODN 2293 SEQ ID NO:2
ODN 2297 SEQ ID NO:2

ODN 2297 SEQ ID NO:2
ODN 2298 SEQ ID NO:2

ODN 2298 SEQ ID NO:2
ODN 2300 SEQ ID NO:3

ODN 2299 SEQ ID NO:2
ODN 2300 SEQ ID NO:2

ODN 2300 SEQ ID NO:2
ODN 2301 SEQ ID NO:2

ODN 2301 SEQ ID NO:2
ODN 2302 SEQ ID NO:3

ODN 2302 SEQ ID NO:2

ODN 2303 SEQ ID NO:2
ODN 2304 SEQ ID NO:3

ODN 2304 SEQ ID NO:2

ODN 2305 SEQ ID NO:2

ODN 2306 SEQ ID NO:3

ODN 2311 SEQ ID NO:3

ODN 2328 SEQ ID NO:3

ODN 2329 SEQ ID NO:3

ODN 2330 SEQ ID NO:3

ODN 2332 SEQ ID NO:3

ODN 2334 SEQ ID NO:3

ODN 2336 SEQ ID NO:3

thioate linkage and each upper

wherein each lower case letter represents phosphorothioate linkage and each upper case letter indicates phosphodiester linkage.

8B7 176. A method of stimulating production of a plurality of type I IFN subtypes, comprising contacting IPCs with an amount of immunostimulatory nucleic acid effective to induce secretion of at least two type I interferons.

5 177. The method of claim 176, wherein the contacting occurs in vivo.

178. The method of claim 176, wherein the contacting occurs in vitro.

10 179. The method of claim 176, wherein the IPCs are precursor type 2 dendritic cells (pDC2s).

180. The method of claim 176, wherein the IPCs are isolated.

15 181. The method of claim 176, wherein the IPCs are induced to secrete at least three type I interferons.

182. The method of claim 176, wherein the IPCs are induced to secrete at least four type I interferons.

20 183. The method of claim 176, wherein the IPCs are induced to secrete at least five type I interferons.

25 184. The method of claim 176, wherein the IPCs are induced to secrete at least six type I interferons.

185. The method of claim 176, wherein the IPCs are induced to secrete at least seven type I interferons.

30 186. The method of claim 176, wherein the IPCs are induced to secrete at least eight type I interferons.

187. The method of claim 176, wherein the immunostimulatory nucleic acid is modified.

09672126 092700

188. The method of claim 176, wherein the immunostimulatory nucleic acid comprises a backbone with at least one nuclease-resistant internucleotide linkage selected from the group consisting of: phosphorothioate, phosphorodithioate, methylphosphonate, and peptide.

189. The method of claim 176, wherein the immunostimulatory nucleic acid comprises at least one nucleotide analog or derivative.

190. The method of claim 176, wherein the immunostimulatory nucleic acid is not a palindrome.

191. The method of claim 176, wherein the immunostimulatory nucleic acid is a CpG nucleic acid.

192. The method of claim 176, wherein the immunostimulatory nucleic acid is a non-CpG nucleic acid.

193. The method of claim 192, wherein the non-CpG immunostimulatory nucleic acid is a T-rich nucleic acid.

194. The method of claim 192, wherein the non-CpG immunostimulatory nucleic acid is a poly-G nucleic acid.

195. The method of claim 176, wherein the immunostimulatory nucleic acid is any combination of at least two nucleic acids selected from the group consisting of: CpG nucleic acids, T-rich nucleic acids, and poly-G nucleic acids.

196. The method of claim 176, wherein the immunostimulatory nucleic acid is between 8 and 100 nucleotides in length.

09672126.092700

197. The method of claim 176, wherein the immunostimulatory nucleic acid is between 12 and 40 nucleotides in length.

198. The method of claim 176, wherein the immunostimulatory nucleic acid has a sequence selected from the group consisting of

5	ggGGTCAACGTTGAggggggG	ODN 1585	SEQ ID NO:1
	tcgtcgttttgcgttttgcgtt	ODN 2022	SEQ ID NO:2
	ggggtcgtcgttttgggggg	ODN 2184	SEQ ID NO:3
	tcgtcgttttgcgttttgggggg	ODN 2185	SEQ ID NO:4
10	ggggtcgacgtcgagggggg	ODN 2192	SEQ ID NO:5
	ggggtcacgtcgagggggg	ODN 2204	SEQ ID NO:6
	ggGGGACGATCGTCggggggG	ODN 2216	SEQ ID NO:7
	gggggtcgtacgacgggggg	ODN 2217	SEQ ID NO:8
	ggGGGACGATATCGTCggggggG	ODN 2245	SEQ ID NO:9
15	ggGGGACGACGTCGTCggggggG	ODN 2246	SEQ ID NO:10
	ggGGGACGAGCTCGTCggggggG	ODN 2247	SEQ ID NO:11
	ggGGGACGTACGTCggggggG	ODN 2248	SEQ ID NO:12
	ggGGGACGATCGTTGggggggG	ODN 2252	SEQ ID NO:13
	ggGGAACGATCGTCggggggG	ODN 2253	SEQ ID NO:14
20	ggGGGGACGATCGTCggggggG	ODN 2254	SEQ ID NO:15
	ggGGGACGATCGTCGggggggG	ODN 2255	SEQ ID NO:16
	ggGGGTCATCGATGAggggggG	ODN 2260	SEQ ID NO:17
	ggGGTCGTCGACGAggggggG	ODN 2293	SEQ ID NO:18
	ggGGTCGTTTCAACGAggggggG	ODN 2294	SEQ ID NO:19
25	ggGGACGTTTCAACGTggggggG	ODN 2295	SEQ ID NO:20
	ggGGAACGACGTCGTTggggggG	ODN 2297	SEQ ID NO:21
	ggGGAACGTACGTCggggggG	ODN 2298	SEQ ID NO:22
	ggGGAACGTACGTACGTTggggggG	ODN 2299	SEQ ID NO:23
	ggGGTCACCGGTGAggggggG	ODN 2300	SEQ ID NO:24
30	ggGGTCGACGTACGTCGAggggggG	ODN 2301	SEQ ID NO:25
	ggGGACCGGTACCGGTggggggG	ODN 2302	SEQ ID NO:26
	ggGTCGACGTCGAggggggG	ODN 2303	SEQ ID NO:27
	ggGGTCGACGTCGaggggg	ODN 2304	SEQ ID NO:28
	ggGGAACGTTAACGTTggggggG	ODN 2305	SEQ ID NO:29
35	ggGGACGTCGACGTggggggG	ODN 2306	SEQ ID NO:30
	ggGGGTCGTTTCGTTggggggG	ODN 2311	SEQ ID NO:31
	ggGACGATCGTCGggggggG	ODN 2328	SEQ ID NO:32
	ggGTCGTCGACGAgggggggG	ODN 2329	SEQ ID NO:33
	ggTCGTCGACGAgggggggG	ODN 2330	SEQ ID NO:34
40	ggGGACGATCGTCGggggggG	ODN 2332	SEQ ID NO:35
	ggGGTCGACGTCGACGTCGAggggggG	ODN 2334	SEQ ID NO:36, and
	ggGGACGACGTCGTGggggggG	ODN 2336	SEQ ID NO:37,

wherein each lower case letter represents phosphorothioate linkage and each upper case letter indicates phosphodiester linkage.

199. A method of inhibiting IL-12 production, comprising
 contacting IL-12-producing cells, in the presence of interferon-producing cells under
 conditions in which the IL-12-producing cells normally produce IL-12, with an
 immunostimulatory nucleic acid in an amount effective for inducing secretion of type I
 interferon.

200. The method of claim 199, wherein the immunostimulatory nucleic acid has a
 sequence selected from the group consisting of

10	ggGGTCAACGTTGAggggggG	ODN 1585	SEQ ID NO:1
	tcgtcgttttgcgttttgcgtt	ODN 2022	SEQ ID NO:2
	ggggtcgtcgttttgggggg	ODN 2184	SEQ ID NO:3
	tcgtcgttttgcgttttgggggg	ODN 2185	SEQ ID NO:4
	ggggtcgacgtcgagggggg	ODN 2192	SEQ ID NO:5
15	ggggtcacgtcgagggggg	ODN 2204	SEQ ID NO:6
	ggGGGACGATCGTCggggggG	ODN 2216	SEQ ID NO:7
	gggggtcgtacgacgggggg	ODN 2217	SEQ ID NO:8
	ggGGGACGATATCGTCggggggG	ODN 2245	SEQ ID NO:9
	ggGGGACGACGTCGTCggggggG	ODN 2246	SEQ ID NO:10
20	ggGGGACGAGCTCGTCggggggG	ODN 2247	SEQ ID NO:11
	ggGGGACGTACGTCggggggG	ODN 2248	SEQ ID NO:12
	ggGGGACGATCGTTggggggG	ODN 2252	SEQ ID NO:13
	ggGGAACGATCGTCggggggG	ODN 2253	SEQ ID NO:14
	ggGGGGACGATCGTCggggggG	ODN 2254	SEQ ID NO:15
25	ggGGGACGATCGTCggggggG	ODN 2255	SEQ ID NO:16
	ggGGGTCATCGATGAggggggG	ODN 2260	SEQ ID NO:17
	ggGGTCGTCGACGAggggggG	ODN 2293	SEQ ID NO:18
	ggGGTCGTTTCAACGAggggggG	ODN 2294	SEQ ID NO:19
	ggGGACGTTTCAACGTggggggG	ODN 2295	SEQ ID NO:20
30	ggGGAACGACGTCGTTggggggG	ODN 2297	SEQ ID NO:21
	ggGGAACGTACGTCggggggG	ODN 2298	SEQ ID NO:22
	ggGGAACGTACGTACGTTggggggG	ODN 2299	SEQ ID NO:23
	ggGGTCACCGGTGAggggggG	ODN 2300	SEQ ID NO:24
	ggGGTCGACGTACGTCGAggggggG	ODN 2301	SEQ ID NO:25
35	ggGGACCGGTACCGGTggggggG	ODN 2302	SEQ ID NO:26
	ggGTCGACGTGAggggggG	ODN 2303	SEQ ID NO:27
	ggGGTCGACGTGAgggggg	ODN 2304	SEQ ID NO:28
	ggGGAACGTTAACGTTggggggG	ODN 2305	SEQ ID NO:29
	ggGGACGTCGACGTggggggG	ODN 2306	SEQ ID NO:30
40	ggGGGTCGTTTCGTTggggggG	ODN 2311	SEQ ID NO:31
	ggGACGATCGTCGggggggG	ODN 2328	SEQ ID NO:32
	ggGTCGTCGACGAgggggggG	ODN 2329	SEQ ID NO:33
	ggTCGTCGACGAGggggggG	ODN 2330	SEQ ID NO:34
	ggGGACGATCGTCGggggggG	ODN 2332	SEQ ID NO:35

002260 " 922700

ggGGTCGACGTCGACGTCGAGggggggG ODN 2334 SEQ ID NO:36, and
ggGGACGACGTCGTGggggggG ODN 2336 SEQ ID NO:37,

wherein each lower case letter represents phosphorothioate linkage and each upper case letter indicates phosphodiester linkage.

5

201. An isolated nucleic acid having a sequence selected from the group consisting of:

tcgtcgttttgcgttttgcgtt	ODN 2022	SEQ ID NO:2
ggggtcgtcgttttgggggg	ODN 2184	SEQ ID NO:3
tcgtcgttttgcgttttgggggg	ODN 2185	SEQ ID NO:4
10 ggggtcgacgtcgagggggg	ODN 2192	SEQ ID NO:5
ggggtcacgtcgagggggg	ODN 2204	SEQ ID NO:6
ggGGGACGATCGTCggggggG	ODN 2216	SEQ ID NO:7
gggggtcgtacgacgggggg	ODN 2217	SEQ ID NO:8
ggGGGACGATATCGTCggggggG	ODN 2245	SEQ ID NO:9
15 ggGGGACGACGTCGTCggggggG	ODN 2246	SEQ ID NO:10
ggGGGACGAGCTCGTCggggggG	ODN 2247	SEQ ID NO:11
ggGGGACGTACGTCggggggG	ODN 2248	SEQ ID NO:12
ggGGGACGATCGTTGggggggG	ODN 2252	SEQ ID NO:13
ggGGAACGATCGTCggggggG	ODN 2253	SEQ ID NO:14
20 ggGGGACGATCGTCggggggG	ODN 2254	SEQ ID NO:15
ggGGGACGATCGTCggggggG	ODN 2255	SEQ ID NO:16
ggGGGTCATCGATGAggggggG	ODN 2260	SEQ ID NO:17
ggGGTCGTCGACGAggggggG	ODN 2293	SEQ ID NO:18
ggGGTCGTTTCAACGAggggggG	ODN 2294	SEQ ID NO:19
25 ggGGACGTTTCAACGTggggggG	ODN 2295	SEQ ID NO:20
ggGGAACGACGTCGTTggggggG	ODN 2297	SEQ ID NO:21
ggGGAACGTACGTCggggggG	ODN 2298	SEQ ID NO:22
ggGGAACGTACGTACGTTggggggG	ODN 2299	SEQ ID NO:23
ggGGTCACCGGTGAggggggG	ODN 2300	SEQ ID NO:24
30 ggGGTCGACGTACGTCGAggggggG	ODN 2301	SEQ ID NO:25
ggGGACCGGTACCGGTggggggG	ODN 2302	SEQ ID NO:26
ggGTCGACGTCGAggggggG	ODN 2303	SEQ ID NO:27
ggGGTCGACGTCGAggggg	ODN 2304	SEQ ID NO:28
ggGGAACGTAAACGTTggggggG	ODN 2305	SEQ ID NO:29
35 ggGGACGTCGACGTgggggG	ODN 2306	SEQ ID NO:30
ggGGGTCGTTTCGTTggggggG	ODN 2311	SEQ ID NO:31
ggGACGATCGTCGggggggG	ODN 2328	SEQ ID NO:32
ggGTCGTCGACGAgggggggG	ODN 2329	SEQ ID NO:33
ggTCGTCGACGAGggggggG	ODN 2330	SEQ ID NO:34
40 ggGGACGATCGTCGggggggG	ODN 2332	SEQ ID NO:35
ggGGTCGACGTCGACGTCGAGggggggG	ODN 2334	SEQ ID NO:36, and
ggGGACGACGTCGTGggggggG	ODN 2336	SEQ ID NO:37,

wherein each lower case letter represents phosphorothioate linkage and each upper case letter indicates phosphodiester linkage.

002260" 9272760

202. A pharmaceutical composition comprising
an isolated nucleic acid having a sequence selected from the group consisting of:

	tcgtcgttttgcgttttgcgtt	ODN 2022	SEQ ID NO:2
5	ggggtcgtcgttttgggggg	ODN 2184	SEQ ID NO:3
	tcgtcgttttgcgttttgggggg	ODN 2185	SEQ ID NO:4
	ggggtcgacgtcgagggggg	ODN 2192	SEQ ID NO:5
	ggggtcacgtcgagggggg	ODN 2204	SEQ ID NO:6
	ggGGGACGATCGTCgggggG	ODN 2216	SEQ ID NO:7
10	gggggtcgtacgacgggggg	ODN 2217	SEQ ID NO:8
	ggGGGACGATATCGTCgggggG	ODN 2245	SEQ ID NO:9
	ggGGGACGACGTCGTCgggggG	ODN 2246	SEQ ID NO:10
	ggGGGACGAGCTCGTCgggggG	ODN 2247	SEQ ID NO:11
	ggGGGACGTACGTCgggggG	ODN 2248	SEQ ID NO:12
15	ggGGGACGATCGTTGggggG	ODN 2252	SEQ ID NO:13
	ggGGAACGATCGTCgggggG	ODN 2253	SEQ ID NO:14
	ggGGGACGATCGTCgggggG	ODN 2254	SEQ ID NO:15
	ggGGGACGATCGTCGgggggG	ODN 2255	SEQ ID NO:16
	ggGGGTCATCGATGAgggggG	ODN 2260	SEQ ID NO:17
20	ggGGTCGTCGACGAgggggG	ODN 2293	SEQ ID NO:18
	ggGGTCGTTTGAACGAgggggG	ODN 2294	SEQ ID NO:19
	ggGGACGTTTGAACGTgggggG	ODN 2295	SEQ ID NO:20
	ggGGAACGACGTCGTTgggggG	ODN 2297	SEQ ID NO:21
	ggGGAACGTACGTCgggggG	ODN 2298	SEQ ID NO:22
25	ggGGAACGTACGTACGTTgggggG	ODN 2299	SEQ ID NO:23
	ggGGTCACCGGTGAgggggG	ODN 2300	SEQ ID NO:24
	ggGGTCGACGTACGTCGAgggggG	ODN 2301	SEQ ID NO:25
	ggGGACCGGTACCGGTgggggG	ODN 2302	SEQ ID NO:26
	ggGTCGACGTCGAgggggG	ODN 2303	SEQ ID NO:27
30	ggGGTCGACGTCGagggg	ODN 2304	SEQ ID NO:28
	ggGGAACGTAAACGTTgggggG	ODN 2305	SEQ ID NO:29
	ggGGACGTCGACGTgggggG	ODN 2306	SEQ ID NO:30
	ggGGGTCGTTTCGTTgggggG	ODN 2311	SEQ ID NO:31
	ggGACGATCGTCGgggggG	ODN 2328	SEQ ID NO:32
35	ggGTCGTCGACGAggggggG	ODN 2329	SEQ ID NO:33
	ggTCGTCGACGAGgggggG	ODN 2330	SEQ ID NO:34
	ggGGACGATCGTCGgggggG	ODN 2332	SEQ ID NO:35
	ggGGTCGACGTCGACGTCGAGgggggG	ODN 2334	SEQ ID NO:36, and
	ggGGACGACGTCGTCGgggggG	ODN 2336	SEQ ID NO:37,

40 wherein each lower case letter represents phosphorothioate linkage and each upper case letter indicates phosphodiester linkage; and
a pharmaceutically acceptable carrier.

203. The pharmaceutical composition of claim 202, further comprising IFN- α .